

Methods for the construction of generators of algebraic curvature tensors

Bernd Fiedler*

Eichelbaumstr. 13, D-04249 Leipzig, Germany. E-mail: bfiedler@femath.de

Dedicated to the memory of Professor Brian G. Wybourne.

Abstract. We demonstrate the use of several tools from Algebraic Combinatorics such as Young tableaux, symmetry operators, the Littlewood-Richardson rule and discrete Fourier transforms of symmetric groups in investigations of algebraic curvature tensors.

In [10, 12, 13] we constructed and investigated generators of algebraic curvature tensors and algebraic covariant derivative curvature tensors. These investigations followed the example of the paper [14] by S.A. Fulling, R.C. King, B.G. Wybourne and C.J. Cummins and applied tools from Algebraic Combinatorics such as Young tableaux, symmetry operators (in particular Young symmetrizers), the Littlewood-Richardson rule, but also discrete Fourier transforms of symmetric groups. The present paper is a short summary of [10, 12, 13] in which we want to demonstrate the use of these methods.

1. The problem

Let V be a finite dimensional $\mathbb{K}$ -vector space, $\mathbb{K} = \mathbb{R}, \mathbb{C}$, and let $\mathcal{T}_r V$ denote the $\mathbb{K}$ -vector space of covariant tensors of order r over V .

DEFINITION 1. *Algebraic curvature tensors* $\mathfrak{R} \in \mathcal{T}_4 V$ and *algebraic covariant derivative curvature tensors* $\mathfrak{R}' \in \mathcal{T}_5 V$ are tensors of order 4 or 5 whose coordinates satisfy

$$\begin{aligned} \mathfrak{R}_{ijkl} &= -\mathfrak{R}_{jikl} = \mathfrak{R}_{klij} & \mathfrak{R}'_{ijklm} &= -\mathfrak{R}'_{jiklm} = \mathfrak{R}'_{klijm} \\ \mathfrak{R}_{ijkl} + \mathfrak{R}_{iklj} + \mathfrak{R}_{iljk} &= 0 & \mathfrak{R}'_{ijklm} + \mathfrak{R}'_{ikljm} + \mathfrak{R}'_{iljkm} &= 0 \\ & & \mathfrak{R}'_{ijklm} + \mathfrak{R}'_{ijlmk} + \mathfrak{R}'_{ijmkl} &= 0. \end{aligned}$$

They are tensors which possess the same symmetry properties as the Riemannian curvature tensor R_{ijkl} and its covariant derivative $R_{ijkl;m}$ of a Levi-Civita connection ∇ .

* 1991 Mathematics Subject Classification: 53B20, 15A72, 05E10, 16D60, 05-04.

The vector space of algebraic curvature tensors $\mathfrak{R} \in \mathcal{T}_4 V$ is spanned by each of the following sets of tensors (P. Gilkey [16, pp.41-44], B. Fiedler¹ [9])

$$\gamma(S)_{ijkl} := S_{il}S_{jk} - S_{ik}S_{jl}, \quad S \text{ symmetric} \quad (1)$$

$$\alpha(A)_{ijkl} := 2A_{ij}A_{kl} + A_{ik}A_{jl} - A_{il}A_{jk}, \quad A \text{ skew-symmetric.} \quad (2)$$

The vector space of algebraic covariant derivative curvature tensors $\mathfrak{R}' \in \mathcal{T}_5 V$ is spanned by the following set of tensors (P. Gilkey [16, p.236], B. Fiedler [10])

$$\hat{\gamma}(S, \hat{S})_{ijkl} := S_{il}\hat{S}_{jk} - S_{jl}\hat{S}_{ik} + S_{jk}\hat{S}_{il} - S_{ik}\hat{S}_{jl}, \quad S, \hat{S} \text{ symmetric.} \quad (3)$$

PROBLEM 2. In the present paper we search for generators of algebraic curvature tensors $\mathfrak{R}$ or algebraic covariant derivative curvature tensors $\mathfrak{R}'$ which can be formed by a suitable symmetry operator from the following types of tensors

$$\mathfrak{R} : U \otimes w, \quad U \in \mathcal{T}_3 V, w \in \mathcal{T}_1 V, \quad (4)$$

$$\mathfrak{R}' : U \otimes W, \quad U \in \mathcal{T}_3 V, W \in \mathcal{T}_2 V, \quad (5)$$

where W and U belong to symmetry classes of $\mathcal{T}_2 V$ and $\mathcal{T}_3 V$ which are defined by minimal right ideals $\mathfrak{r} \subset \mathbb{K}[\mathcal{S}_2]$ and $\hat{\mathfrak{r}} \subset \mathbb{K}[\mathcal{S}_3]$, respectively.

We use Boerner's definition [1, p.127] of symmetry classes of tensors. An element $a = \sum_{p \in \mathcal{S}_r} a(p)p \in \mathbb{K}[\mathcal{S}_r]$ of the *group ring* of the *symmetric group* $\mathcal{S}_r$ can be considered a *symmetry operator* for covariant tensors $T \in \mathcal{T}_r V$. The action of a on T is defined by:

$$(aT)(v_1, \dots, v_r) := \sum_{p \in \mathcal{S}_r} a(p) T(v_{p(1)}, \dots, v_{p(r)}) \quad , \quad v_i \in V. \quad (6)$$

DEFINITION 3. Let $\mathfrak{r} \subseteq \mathbb{K}[\mathcal{S}_r]$ be a right ideal of $\mathbb{K}[\mathcal{S}_r]$ for which an $a \in \mathfrak{r}$ and a $T \in \mathcal{T}_r V$ exist such that $aT \neq 0$. Then the tensor set

$$\mathcal{T}_{\mathfrak{r}} := \{aT \mid a \in \mathfrak{r}, T \in \mathcal{T}_r V\} \quad (7)$$

is called the *symmetry class* of tensors defined by $\mathfrak{r}$.

Boerner [1, p.127] showed: If $e \in \mathbb{K}[\mathcal{S}_r]$ is a generating idempotent of $\mathfrak{r}$, i.e. $\mathfrak{r} = e \cdot \mathbb{K}[\mathcal{S}_r]$, then it holds

$$\boxed{T \in \mathcal{T}_r V \text{ belongs to } \mathcal{T}_{\mathfrak{r}} \Leftrightarrow eT = T.}$$

¹[9] uses also tools of Algebraic Combinatorics, in particular plethysms.

2. Young symmetrizers

Young symmetrizers are important symmetry operators. In particular the symmetries of the Riemann tensor R and its covariant derivatives are characterized by a Young symmetrizer. First we define Young tableaux.

A *Young tableau* t of $r \in \mathbb{N}$ is an arrangement of r boxes such that

1. the numbers λ_i of boxes in the rows $i = 1, \dots, l$ form a decreasing sequence $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_l > 0$ with $\lambda_1 + \dots + \lambda_l = r$,
2. the boxes are fulfilled by the numbers $1, 2, \dots, r$ in any order.

For instance, the following graphics shows a Young tableau of $r = 16$.

$$\left. \begin{array}{l} \lambda_1 = 5 \\ \lambda_2 = 4 \\ \lambda_3 = 4 \\ \lambda_4 = 2 \\ \lambda_5 = 1 \end{array} \right\} \begin{array}{|c|c|c|c|c|} \hline 11 & 2 & 5 & 4 & 12 \\ \hline 9 & 6 & 16 & 15 & \\ \hline 8 & 14 & 1 & 7 & \\ \hline 13 & 3 & & & \\ \hline 10 & & & & \\ \hline \end{array} \right\} = t.$$

Obviously, the unfilled arrangement of boxes, the *Young frame*, is characterized by a partition $\lambda = (\lambda_1, \dots, \lambda_l) \vdash r$ of r .

If a Young tableau t of a partition $\lambda \vdash r$ is given, then the *Young symmetrizer* y_t of t is defined by

$$y_t := \sum_{p \in \mathcal{H}_t} \sum_{q \in \mathcal{V}_t} \text{sign}(q) p \circ q \quad (8)$$

where $\mathcal{H}_t$, $\mathcal{V}_t$ are the groups of the *horizontal* or *vertical permutations* of t which only permute numbers within rows or columns of t , respectively. The Young symmetrizers of $\mathbb{K}[\mathcal{S}_r]$ are essentially idempotent and define decompositions

$$\mathbb{K}[\mathcal{S}_r] = \bigoplus_{\lambda \vdash r} \bigoplus_{t \in \mathcal{ST}_\lambda} \mathbb{K}[\mathcal{S}_r] \cdot y_t, \quad \mathbb{K}[\mathcal{S}_r] = \bigoplus_{\lambda \vdash r} \bigoplus_{t \in \mathcal{ST}_\lambda} y_t \cdot \mathbb{K}[\mathcal{S}_r] \quad (9)$$

of $\mathbb{K}[\mathcal{S}_r]$ into minimal left or right ideals $\mathbb{K}[\mathcal{S}_r] \cdot y_t$, $y_t \cdot \mathbb{K}[\mathcal{S}_r]$. In (9), the symbol $\mathcal{ST}_\lambda$ denotes the set of all standard tableaux of the partition λ . Standard tableaux are Young tableaux in which the entries of every row and every column form an increasing number sequence.¹

The inner sums of (9) are minimal two-sided ideals

$$\mathfrak{a}_\lambda := \bigoplus_{t \in \mathcal{ST}_\lambda} \mathbb{K}[\mathcal{S}_r] \cdot y_t = \bigoplus_{t \in \mathcal{ST}_\lambda} y_t \cdot \mathbb{K}[\mathcal{S}_r] \quad (10)$$

of $\mathbb{K}[\mathcal{S}_r]$. The set of all Young symmetrizers y_t which lie in $\mathfrak{a}_\lambda$ is equal to the set of all y_t whose tableau t has the frame $\lambda \vdash r$. Furthermore two minimal left ideals

¹About Young symmetrizers and Young tableaux see for instance [1, 14, 15, 17, 18, 20, 21].

$\mathfrak{l}_1, \mathfrak{l}_2 \subseteq \mathbb{K}[\mathcal{S}_r]$ or two minimal right ideals $\mathfrak{r}_1, \mathfrak{r}_2 \subseteq \mathbb{K}[\mathcal{S}_r]$ are *equivalent* iff they lie in the same ideal $\mathfrak{a}_\lambda$. Now we say that a symmetry class $\mathcal{T}_\mathfrak{r}$ *belongs to* $\lambda \vdash r$ iff $\mathfrak{r} \subseteq \mathfrak{a}_\lambda$.

S.A. Fulling, R.C. King, B.G. Wybourne and C.J. Cummins showed in [14] that the symmetry classes of the Riemann tensor R and its *symmetrized² covariant derivatives*

$$\left(\nabla^{(u)} R\right)_{ijkl s_1 \dots s_u} := \nabla_{(s_1} \nabla_{s_2} \dots \nabla_{s_u)} R_{ijkl} = R_{ijkl; (s_1 \dots s_u)} \quad (11)$$

are generated by special Young symmetrizers³.

PROPOSITION 4. (Fulling, King, Wybourne, Cummins)

Let ∇ be the Levi-Civita connection of a pseudo-Riemannian metric g . For $u \geq 0$ the symmetrized covariant derivatives $\nabla^{(u)} R$ fulfil

$$e_t^* \nabla^{(u)} R = \nabla^{(u)} R \quad (12)$$

where $e_t := y_t(u+1)/(2 \cdot (u+3)!)$ is an idempotent which is formed from the Young symmetrizer y_t of the standard tableau

$$t = \begin{array}{|c|c|c|c|c|} \hline 1 & 3 & 5 & \dots & \dots \\ \hline 2 & 4 & & & \\ \hline \end{array} \begin{array}{|c|} \hline (u+4) \\ \hline \end{array}. \quad (13)$$

The $'*$ ' in (12) is the mapping $*$: $a = \sum_{p \in \mathcal{S}_r} a(p) p \mapsto a^* := \sum_{p \in \mathcal{S}_r} a(p) p^{-1}$.

We see from Proposition 4 that the tensor fields $\nabla^{(u)} R$ belong to the symmetry class which is defined by the symmetrizer y_t^* of (13), more precisely, by the right ideal $\mathfrak{r} = y_t^* \cdot \mathbb{K}[\mathcal{S}_{u+4}]$. In the special case of algebraic tensors $\mathfrak{R}, \mathfrak{R}'$ we have the following corollary (see [10]):

COROLLARY 5. *Let us denote by t and t' the standard tableaux*

$$t = \begin{array}{|c|c|} \hline 1 & 3 \\ \hline 2 & 4 \\ \hline \end{array}, \quad t' = \begin{array}{|c|c|c|} \hline 1 & 3 & 5 \\ \hline 2 & 4 & \\ \hline \end{array}. \quad (14)$$

Then a tensor $T \in \mathcal{T}_4 V$ or $\tilde{T} \in \mathcal{T}_5 V$ is an algebraic curvature tensor or an algebraic covariant derivative curvature tensor iff these tensors satisfy

$$y_t^* T = 12 T, \quad y_{t'}^* \tilde{T} = 24 \tilde{T}, \quad (15)$$

respectively. Thus the symmetry classes of the algebraic tensors $\mathfrak{R}, \mathfrak{R}'$ are defined by the minimal right ideals $y_t^ \cdot \mathbb{K}[\mathcal{S}_4], y_{t'}^* \cdot \mathbb{K}[\mathcal{S}_5]$ which belong to the partitions $(2, 2) \vdash 4, (3, 2) \vdash 5$.*

²(...) denotes the symmetrization with respect to the indices $s_1, \dots, s_u$.

³A proof of this result of [14] can be found in [5, Sec.6], too. See also [10] for more details.

3. Symmetry classes of $\mathcal{T}_3V$ belonging to $\lambda = (21)$

The group ring $\mathbb{K}[\mathcal{S}_3]$ contains the minimal 2-sided ideals $\mathfrak{a}_{(3)}, \mathfrak{a}_{(21)}, \mathfrak{a}_{(13)}$. The 2-sided ideals $\mathfrak{a}_{(3)}, \mathfrak{a}_{(13)} \subset \mathbb{K}[\mathcal{S}_3]$ have dimension¹ 1 and define consequently unique symmetry classes of $\mathcal{T}_3V$. The 2-sided ideal $\mathfrak{a}_{(21)} \subset \mathbb{K}[\mathcal{S}_3]$, however, has dimension 4 and contains an infinite set of minimal right ideals $\hat{\mathfrak{r}}$ (of dimension 2) which lead to an infinite set of possible symmetry classes for the tensor $U \in \mathcal{T}_3V$.

We use discrete Fourier transforms to determine a generating idempotent for every such $\hat{\mathfrak{r}}$.

DEFINITION 6. A *discrete Fourier transform*² for $\mathcal{S}_r$ is an isomorphism

$$D : \mathbb{K}[\mathcal{S}_r] \rightarrow \bigotimes_{\lambda \vdash r} \mathbb{K}^{d_\lambda \times d_\lambda}, \quad \sum_{p \in \mathcal{S}_r} a(p) p \mapsto \begin{pmatrix} A_{\lambda_1} & & 0 \\ & A_{\lambda_2} & \\ 0 & & A_{\lambda_k} \end{pmatrix} \quad (16)$$

according to Wedderburn's theorem which maps the group ring $\mathbb{K}[\mathcal{S}_r]$ onto an outer direct product $\bigotimes_{\lambda \vdash r} \mathbb{K}^{d_\lambda \times d_\lambda}$ of full matrix rings $\mathbb{K}^{d_\lambda \times d_\lambda}$.

In (16) the matrix ring $\mathbb{K}^{d_\lambda \times d_\lambda}$ corresponds to the minimal two-sided ideal $\mathfrak{a}_\lambda$ of $\mathbb{K}[\mathcal{S}_r]$. For $\mathcal{S}_3$ we have a mapping

$$D : a = \sum_{p \in \mathcal{S}_3} a(p) p \mapsto \begin{pmatrix} A_{(3)} & & 0 \\ & A_{(21)} & \\ 0 & & A_{(13)} \end{pmatrix}, \quad (17)$$

where $A_{(3)}$ and $A_{(13)}$ are 1×1 -matrices and $A_{(21)}$ is a 2×2 -matrix. It holds $a \in \mathfrak{a}_{(21)}$ iff $A_{(3)} = A_{(13)} = 0$. In [10] we proved

PROPOSITION 7. *Every minimal right ideal $\mathfrak{r} \subset \mathbb{K}^{2 \times 2}$ is generated by exactly one of the following (primitive) idempotents*

$$X_\infty := \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, \quad X_\nu := \begin{pmatrix} 1 & 0 \\ \nu & 0 \end{pmatrix}, \quad \nu \in \mathbb{K}. \quad (18)$$

From (18) we obtain the generating idempotents for the right ideals $\hat{\mathfrak{r}} \subset \mathfrak{a}_{(21)} \subset \mathbb{K}[\mathcal{S}_3]$ by

$$\xi_\nu = D^{-1} \begin{pmatrix} 0 & & \\ & X_\nu & \\ & & 0 \end{pmatrix} = \begin{cases} \frac{1}{3} \{ [1, 2, 3] - [2, 1, 3] - [2, 3, 1] + [3, 2, 1] \}, & \nu = \infty \\ \frac{1}{3} \{ [1, 2, 3] + \nu [1, 3, 2] + (1 - \nu) [2, 1, 3] \\ - \nu [2, 3, 1] + (-1 + \nu) [3, 1, 2] - [3, 2, 1] \}, & \text{else.} \end{cases} \quad (19)$$

¹The dimension of a minimal left or right ideal can be calculated from the Young frame belonging to it by the hook length formula (see e.g. [1, 14, 17, 18].)

²Three discrete Fourier transforms are known for symmetric groups $\mathcal{S}_r$: *Young's natural representation* [1, pp.102-108], *Young's seminormal representation* [2], [3, p.130], [18, p.76] and *Young's orthogonal representation* [2, 3, 18] of $\mathcal{S}_r$.

Note that the above connection between ν and ξ_ν depends on the concrete discrete Fourier transform D which is used in (19). The above formula (19) was determined by means of the *Mathematica* package PERMS [7] whose discrete Fourier transform D is based on *Young's natural representation* of $\mathcal{S}_r$.

Examples of tensor fields with a (21)-symmetry can be constructed from covariant derivatives of symmetric or alternating 2-fields.

PROPOSITION 8. ³ *Let ∇ be a torsion-free covariant derivative on a C^∞ -manifold M , $\dim M \geq 2$. Further let $\psi, \omega \in \mathcal{T}_2 M$ be differentiable tensor fields of order 2 which are symmetric or skew-symmetric, respectively. Then for every point $p \in M$ the tensors*

$$(\nabla\psi - \text{sym}(\nabla\psi))|_p \Rightarrow \nu = 0 \quad , \quad (\nabla\omega - \text{alt}(\nabla\omega))|_p \Rightarrow \nu = 2 \quad (20)$$

lie in (21)-symmetry classes whose generating idempotents ξ_ν belong to the above ν -values 0 or 2. ('sym' = symmetrization, 'alt' = anti-symmetrization.)

In a future paper we will show that tensor fields with a (21)-symmetry also occur in curvature formulas connected with *stationary* and *static space-times*.

4. Our main results

Now we formulate our main results which were proved in [10, 13]. The next section will give some ideas of the proofs.

THEOREM 9. *A solution of Problem 2 can be constructed at most from such products (4) or (5) whose factors belong to the following symmetry classes:*

product		partitions of the symm. classes	
$\mathfrak{R} : U \otimes w$	(a)	$U \rightarrow (21)$	
$\mathfrak{R}' : U \otimes W$	(a')	$U \rightarrow (3) \quad , \quad W \rightarrow (2)$	U and W symmetric
	(b')	$U \rightarrow (21) \quad , \quad W \rightarrow (2)$	W symmetric
	(c')	$U \rightarrow (21) \quad , \quad W \rightarrow (1^2)$	W skew-symmetric

The case (a') of Theorem 9 is realized in formula (3). The cases (a), (b') and (c') of Theorem 9 lead to

THEOREM 10. *The products (a), (b'), (c') lead to generators*

$$y_t^*(U \otimes w) \quad , \quad y_{t'}^*(U \otimes S) \quad , \quad y_{t'}^*(U \otimes A)$$

of the spaces of algebraic tensors $\mathfrak{R}, \mathfrak{R}'$ if and only if the generating idempotent ξ_ν of the symmetry class of U fulfills

$$\boxed{\nu \neq \frac{1}{2}}.$$

Here t and t' are the Young tableaux (14).

³See [11].

5. Ideas of the proofs

5.1. USE OF THE LITTLEWOOD-RICHARDSON RULE (PROOF OF THEOREM 9)

When we consider a right ideal $\mathfrak{r}$ that defines the symmetry class of a product $T_1 \otimes T_2$ of tensors of order r_1, r_2 , then we can determine information about the decomposition of $\mathfrak{r}$ into minimal right ideals by means of *Littlewood-Richardson products* (see [20, 14, 6, 8, 10]). Let $\mathfrak{r}_1, \mathfrak{r}_2$ be the right ideals defining the symmetry classes of T_1, T_2 . We consider the left ideals $\mathfrak{l}_i := \mathfrak{r}_i^*$ representation spaces of subrepresentations α_i of the natural representation of $\mathcal{S}_{r_i}$. Then we have $\mathfrak{r} = \mathfrak{l}^*$ where the left ideal $\mathfrak{l}$ is the representation space of the Littlewood-Richardson product $\alpha_1 \alpha_2 := (\alpha_1 \# \alpha_2) \uparrow \mathcal{S}_{r_1+r_2}$. (' $\#$ ' denotes the outer tensor product and ' $\uparrow$ ' the forming of the induced representation.)

For the tensor products (4), (5) we have to calculate the following *Littlewood-Richardson products* by means of the *Littlewood-Richardson rule*¹:

$$\begin{array}{ll} \mathfrak{R} : & \begin{array}{l} [3][1] \sim [4] + [3\ 1] \\ [2\ 1][1] \sim [3\ 1] + [2^2] + [2\ 1^2] \\ [1^3][1] \sim [2\ 1^2] + [1^4] \end{array} & \mathfrak{R}' : & \begin{array}{l} [3][2] \sim [5] + [3\ 2] + [4\ 1] \\ [3][1^2] \sim [4\ 1] + [3\ 1^2] \\ [2\ 1][2] \sim [3\ 2] + [4\ 1] + [2^2\ 1] + [3\ 1^2] \\ [2\ 1][1^2] \sim [3\ 2] + [2^2\ 1] + [3\ 1^2] + [2\ 1^3] \\ [1^3][2] \sim [3\ 1^2] + [2\ 1^3] \\ [1^3][1^2] \sim [2^2\ 1] + [2\ 1^3] + [1^5] \end{array} \end{array}$$

Only the products $[2\ 1][1]$ for $\mathfrak{R}$ and $[3][2], [2\ 1][2], [2\ 1][1^2]$ for $\mathfrak{R}'$ contain minimal right ideals that belong to the partitions (2 2) for $\mathfrak{R}$ and (3 2) for $\mathfrak{R}'$. We will definitely obtain $y_t^*(U \otimes w) = 0$ or $y_{t'}^*(U \otimes W) = 0$ if the ideal $\mathfrak{r}$ or $\mathfrak{r}'$ of $W \otimes w$ or $U \otimes W$ does not possess a subideal belonging to (2 2) or (3 2), since then $y_t^* \in \mathfrak{a}_{(2\ 2)}, y_{t'}^* \in \mathfrak{a}'_{(3\ 2)}$ but $\mathfrak{r} \cap \mathfrak{a}_{(2\ 2)} = 0, \mathfrak{r}' \cap \mathfrak{a}'_{(3\ 2)} = 0$.

5.2. A STEP OF THE PROOF OF THEOREM 10

Let us consider the example of expressions $y_{t'}^*(U \otimes S)$ and $y_{t'}^*(U \otimes A)$. To treat such expressions we form the following group ring elements of $\mathbb{K}[\mathcal{S}_5]$:

$$\sigma_{\nu, \epsilon} := y_{t'}^* \cdot \xi'_\nu \cdot \zeta''_\epsilon \quad (21)$$

$$\zeta''_\epsilon := \text{id} + \epsilon(4\ 5) \quad , \quad \epsilon \in \{1, -1\} \quad (22)$$

$$\xi_\nu \mapsto \xi'_\nu \in \mathbb{K}[\mathcal{S}_5] \quad (23)$$

Formula (23) denotes the embedding of the group ring elements $\xi_\nu \in \mathbb{K}[\mathcal{S}_3]$ into $\mathbb{K}[\mathcal{S}_5]$ which is induced by the mapping $\mathcal{S}_3 \rightarrow \mathcal{S}_5$, $[i_1, i_2, i_3] \mapsto [i_1, i_2, i_3, 4, 5]$. The symmetry operator $\xi'_\nu \cdot \zeta''_\epsilon$ maps arbitrary product tensors $T''' \otimes T''$ to products $U \otimes S$ or $U \otimes A$. Using our Mathematica package PERMS [7] we verified $\sigma_{\nu, \epsilon} \neq 0 \Leftrightarrow \nu \neq \frac{1}{2}$. The value $\nu = \frac{1}{2}$ has to be excluded since $\sigma_{\nu, \epsilon} = 0$ and $\mathfrak{R}' = 0$ in this case.

¹See [20, 14, 6].

6. Shortest formulas for generators of $\mathfrak{R}$ and $\mathfrak{R}'$

In this section we want to construct generators (4), (5) whose coordinate representation has a minimal number of summands. To this end we determine systems of linear identities which are satisfied by the coordinates of all tensors from the symmetry class of U . In [6, Sec.III.4.1] we proved

PROPOSITION 11. *Let $\mathfrak{r} \subset \mathbb{K}[\mathcal{S}_r]$ be a d -dimensional right ideal that defines a symmetry class $\mathcal{T}_{\mathfrak{r}}$ of tensors $T \in \mathcal{T}_r V$. If a basis $\{h_1, \dots, h_d\}$ of the left ideal $\mathfrak{l} = \mathfrak{r}^*$ is known, then every solution x_p of the linear $(d \times r!)$ -equation system*

$$\sum_{p \in \mathcal{S}_r} h_i(p) x_p = 0 \quad (i = 1, \dots, d). \quad (24)$$

yields the coefficients for a linear identity

$$\sum_{p \in \mathcal{S}_r} x_p T_{i_{p(1)} \dots i_{p(r)}} = 0 \quad (25)$$

fulfilled by the coordinates of all $T \in \mathcal{T}_{\mathfrak{r}}$.

For our tensors U the rank of the equation system (24) is equal to $\dim \mathfrak{r} = 2$. The columns of (24) are numbered by the permutations $p \in \mathcal{S}_3$.

Now we determine identities (25) for U by the following procedure. We form the system (24) from the idempotent ξ_ν by the determination of a basis¹ from the generating set $\{p \cdot \xi_\nu^* \mid p \in \mathcal{S}_3\}$ of $\mathfrak{l} = \mathfrak{r}^*$. Then for every subset $\mathcal{P} = \{p_1, p_2\} \subset \mathcal{S}_3$ we check the determinant $\Delta_{\mathcal{P}}$ of the corresponding (2×2) -submatrix of (24). If $\Delta_{\mathcal{P}} \neq 0$, then we determine identities (25) of the special form

$$0 = \sum_{p \in \mathcal{P}} x_p^{(\bar{p})} U_{i_{p(1)} i_{p(2)} i_{p(3)}} + U_{i_{\bar{p}(1)} i_{\bar{p}(2)} i_{\bar{p}(3)}} \quad (\bar{p} \in \mathcal{S}_3 \setminus \mathcal{P}). \quad (26)$$

For instance, the set $\mathcal{P} = \{[1, 2, 3], [1, 3, 2]\}$ leads to the determinant $\Delta_{\mathcal{P}}(\nu) = \frac{4}{27}(1 - \nu)(1 + \nu)$ which has the roots $\nu_1 = 1$ and $\nu_2 = -1$. For $\nu \notin \{1, -1\}$ we obtain the identities

$$\begin{aligned} -\frac{\nu^2 - \nu + 1}{\nu^2 - 1} U_{ijk} + \frac{2\nu - 1}{\nu^2 - 1} U_{ikj} + U_{kji} &= 0 \\ \frac{\nu^2 - 2\nu}{\nu^2 - 1} U_{ijk} + \frac{\nu^2 - \nu + 1}{\nu^2 - 1} U_{ikj} + U_{kij} &= 0 \\ \frac{2\nu - 1}{\nu^2 - 1} U_{ijk} - \frac{\nu^2 - \nu + 1}{\nu^2 - 1} U_{ikj} + U_{jki} &= 0 \\ \frac{\nu^2 - \nu + 1}{\nu^2 - 1} U_{ijk} + \frac{\nu^2 - 2\nu}{\nu^2 - 1} U_{ikj} + U_{jik} &= 0 \end{aligned} \quad (27)$$

There exist 15 subsets $\mathcal{P} = \{p_1, p_2\} \subset \mathcal{S}_3$ and consequently 15 systems (27) for U .

¹Faster algorithms which determine a basis also for a large $\mathcal{S}_r$ by means of discrete Fourier transforms were developed in [6].

Now let us consider the coordinates $\frac{1}{24}(y_{\nu'}^*(U \otimes S))_{ijklr}$ of generators for $\mathfrak{A}'$. If we use (27) to express all coordinates of U by U_{ijk} and U_{ikj} we obtain the following sum of 16 terms.

$$\begin{aligned}
& \frac{-(-1+2\nu)}{24(-1+\nu)(1+\nu)} U_{jlr} S_{ik} + \frac{\nu(-1+2\nu)}{24(-1+\nu)(1+\nu)} U_{jrl} S_{ik} + \\
& \frac{-1+2\nu}{24(-1+\nu)(1+\nu)} U_{jkr} S_{il} - \frac{\nu(-1+2\nu)}{24(-1+\nu)(1+\nu)} U_{jrk} S_{il} + \\
& \frac{-1+2\nu}{24(-1+\nu)} U_{jkl} S_{ir} - \frac{-1+2\nu}{24(-1+\nu)} U_{jlk} S_{ir} + \\
& \frac{-1+2\nu}{24(-1+\nu)(1+\nu)} U_{ilr} S_{jk} - \frac{\nu(-1+2\nu)}{24(-1+\nu)(1+\nu)} U_{irl} S_{jk} - \\
& \frac{-1+2\nu}{24(-1+\nu)(1+\nu)} U_{ikr} S_{jl} + \frac{\nu(-1+2\nu)}{24(-1+\nu)(1+\nu)} U_{irk} S_{jl} - \\
& \frac{-1+2\nu}{24(-1+\nu)} U_{ikl} S_{jr} + \frac{-1+2\nu}{24(-1+\nu)} U_{ilk} S_{jr} - \\
& \frac{(-1+2\nu)^2}{24(-1+\nu)(1+\nu)} U_{ijl} S_{kr} - \frac{(-2+\nu)(-1+2\nu)}{24(-1+\nu)(1+\nu)} U_{ilj} S_{kr} + \\
& \frac{(-1+2\nu)^2}{24(-1+\nu)(1+\nu)} U_{ijk} S_{lr} + \frac{(-2+\nu)(-1+2\nu)}{24(-1+\nu)(1+\nu)} U_{ikj} S_{lr}
\end{aligned}$$

This sum has the structure

$$\mathfrak{P}_{i_1 \dots i_5}^{\text{red}} = \sum_{q \in \mathcal{S}_5} \frac{P_q^{\mathcal{P}}(\nu)}{Q_q^{\mathcal{P}}(\nu)} U_{i_{q(1)} i_{q(2)} i_{q(3)}} S_{i_{q(4)} i_{q(5)}}. \quad (28)$$

where $P_q^{\mathcal{P}}(\nu)$ and $Q_q^{\mathcal{P}}(\nu)$ are polynomials. If we determine the set $N_{\mathcal{P}}$ of all roots $\nu \neq \frac{1}{2}$ of the $P_q^{\mathcal{P}}(\nu)$, for which $\Delta_{\mathcal{P}}(\nu) \neq 0$, and set the $\nu \in N_{\mathcal{P}}$ into (28), the length of (28) will decrease.

We determine the minimal length of (28) by this procedure for $y_t^*(U \otimes w)$, $y_{\nu'}^*(U \otimes S)$, $y_{\nu'}^*(U \otimes A)$ and for every of the 15 identity systems of type (27) in the case $\nu \neq \infty$. Table 1 shows the results for $y_{\nu'}^*(U \otimes S)$, $y_{\nu'}^*(U \otimes A)$. (In the column for $\mathcal{P} = \{p_1, p_2\}$ the p_i are denoted by their numbers in the lexicographically ordered $\mathcal{S}_3$.) Furthermore we calculate the lengths of (28) for the 15 systems (27) in the case $\nu = \infty$. Altogether, the number of calculations comes to

$$(3 \text{ generator types}) \times (2 \text{ } \nu\text{-cases}) \times (15 \text{ systems (27)}) = 75.$$

We obtain (see [12, 13])

THEOREM 12. *Let $\dim V \geq 3$. Then the coordinates of $y_t^*(U \otimes w)$, $y_{\nu'}^*(U \otimes S)$, $y_{\nu'}^*(U \otimes A)$ are sums of the following lengths*

		$y_t^*(U \otimes w)$	$y_{\nu'}^*(U \otimes S)$	$y_{\nu'}^*(U \otimes A)$
(a)	generic case for ν	8	16	20
(b)	ν producing minimal length	4	12	10

The computer calculations were carried out by means of the Mathematica packages Ricci [19] and PERMS [7]. Notebooks of the calculations are available on the web page [4].

It is very remarkable that U admits an index commutation symmetry if the coordinates of $y_t^*(U \otimes w)$, $y_{\nu'}^*(U \otimes S)$, $y_{\nu'}^*(U \otimes A)$ have the minimal lengths of case (b) in Theorem 12 (see [12, 13]).

$\mathcal{P}$	$y_{t'}^*(U \otimes S)$		$y_{t'}^*(U \otimes A)$	
	roots of $P_q^{\mathcal{P}}(\nu)$ with $\Delta_{\mathcal{P}}(\nu) \neq 0, \nu \neq 1/2$	length of $\mathfrak{P}_{i_1 \dots i_5}^{\text{red}}$	roots of $P_q^{\mathcal{P}}(\nu)$ with $\Delta_{\mathcal{P}}(\nu) \neq 0, \nu \neq 1/2$	length of $\mathfrak{P}_{i_1 \dots i_5}^{\text{red}}$
12	0	12		
	2	14	2	12
13	-1	14	-1	18
14	-1	14	-1	18
	0	12		
	2	12	2	10
15	-1	12	-1	10
	0	12		
	1	12		
	2	14	2	12
16	-1	12	-1	10
	0	12		
	1	12		
	2	12	2	10
23	-1	14	-1	18
	0	12		
	2	14	2	12
24	-1	14	-1	18
	2	14	2	18
25	-1	12	-1	10
	1	12		
26	-1	12	-1	10
	1	12		
	2	14	2	18
34	0	12		
	2	12	2	10
35	-1	14	-1	12
	0	12		
	1	12		
	2	14	2	12
36	-1	14	-1	12
	0	12		
	1	12		
	2	12	2	10
45	-1	14	-1	12
	1	12		
	2	14	2	18
46	-1	14	-1	12
	1	12		
56	2	14	2	18

TABLE 1. The lengths of $\mathfrak{P}_{i_1 \dots i_5}^{\text{red}}$ for $y_{t'}^*(U \otimes S)$, $y_{t'}^*(U \otimes A)$ and $\nu \neq \infty$.

Bibliography

1. H. Boerner, *Darstellungen von Gruppen*, Springer, Berlin, Göttingen, Heidelberg, 1955.
2. H. Boerner, *Representations of Groups*, 2. revised ed., North-Holland, Amsterdam, 1970.
3. M. Clausen and U. Baum, *Fast Fourier Transforms*, BI Wissenschaftsverlag, Mannheim, Leipzig, Wien, Zürich, 1993.
4. B. Fiedler, *Examples of calculations by means of PERMS*, Mathematica notebooks, Internet <http://www.fiemath.de/pnbks.htm>.
5. B. Fiedler, *Z. Anal. Anw.* **17**(1), 135 (1998).
6. B. Fiedler, *An Algorithm for the Decomposition of Ideals of Semi-Simple Rings and its Application to Symbolic Tensor Calculations by Computer*, Habilitationsschrift, Fakultät für Mathematik und Informatik, Univ. Leipzig, Germany, November 1999.
7. B. Fiedler, *PERMS 2.1 (15.1.1999)*. Mathematisches Institut, Univ. Leipzig, Germany, 1999. Will be sent in to MathSource, Wolfram Research Inc.
8. B. Fiedler, Séminaire Lotharingien de Combinatoire, **B45g**, 16 pp. (2001). El. published: <http://www.mat.univie.ac.at/~slc>, Archive: <http://arXiv.org/abs/math.CO/0211156>.
9. B. Fiedler, Séminaire Lotharingien de Combinatoire, **B48d**, 20 pp. (2002). El. published: <http://www.mat.univie.ac.at/~slc>, Preprint: <http://arXiv.org/abs/math.CO/0212278>.
10. B. Fiedler, *Generators of algebraic covariant derivative curvature tensors and Young symmetrizers*, Preprint: <http://arXiv.org/abs/math.CO/0310020>, 18pp. 2003. Chapter for a book "Progress in Computer Science Research", in prep. by Nova Science Publishers, Inc.
11. B. Fiedler, Séminaire Lotharingien de Combinatoire, **B49f**, 22 pp. (2003). El. published: <http://www.mat.univie.ac.at/~slc>, Preprint: <http://arXiv.org/abs/math.CO/0301042>.
12. B. Fiedler, *Short formulas for algebraic covariant derivative curvature tensors via algebraic combinatorics*, Preprint: <http://arXiv.org/abs/math.CO/0312171>, 38pp. 2003. Paper for Proc. of Conf. "PDEs, Submanifolds and Affine Differential Geometry", TU Berlin and Banach Center, Bedlewo 2003. Submitted.
13. B. Fiedler, *Generators of algebraic curvature tensors based on a (2 1)-symmetry*, Preprint: <http://arxiv.org/abs/math.DG/0411056>, 16 pp. 2004.
14. S. A. Fulling, R. C. King, B. G. Wybourne and C. J. Cummins, *Class. Quantum Grav.* **9**, 1151 (1992).
15. W. Fulton, *Young Tableaux*, Cambridge University Press, Cambridge, New York, Melbourne, 1997.
16. P. B. Gilkey, *Geometric Properties of Natural Operators Defined by the Riemann Curvature Tensor*, World Scientific Publishing Co., Singapore, New Jersey, London, Hong Kong, 2001.
17. G. D. James and A. Kerber, *The Representation Theory of the Symmetric Group*, Addison-Wesley, Reading, Mass., London, Amsterdam, Don Mills, Ont., Sidney, Tokyo, 1981.
18. A. Kerber, *Representations of Permutation Groups*, Lecture Notes in Mathematics vol. 240, 495, Springer-Verlag, Berlin, Heidelberg, New York, 1971, 1975.
19. J. M. Lee, D. Lear, J. Roth, J. Coskey and L. Nave, *Ricci. A Mathematica package for doing tensor calculations in differential geometry. User's Manual. Version 1.32*, Department of Mathematics, Box 354350, University of Washington, Seattle, WA 98195-4350, 1992 - 1998. Ricci's home page: <http://www.math.washington.edu/~lee/Ricci/>.
20. D. E. Littlewood, *The Theory of Group Characters and Matrix Representations of Groups*, 2. ed., Clarendon Press, Oxford, 1950.
21. I. G. Macdonald, *Symmetric Functions and Hall Polynomials*, Clarendon Press, Oxford, 1979.